arXiv:1209.1747v1 [hep-ph] 8 Sep 2012

Correlation femtoscopy of small systems

Yu.M. Sinyukov and V.M. Shapoval
Bogolyubov Institute for Theoretical Physics,
Metrolohichna str. 14b, 03680 Kiev, Ukraine

Abstract

The basic principles of the correlation femtoscopy, including its correspondence to the Hanbury Brown and Twiss intensity interferometry, are re-examined. The main subject of the paper is an analysis of the correlation femtoscopy of small systems with typical sizes about 1 fm. The uncertainty principle plays an important role for correlation femtoscopy of such systems. It is taken into account by the use of the formalism of partially coherent phases in the amplitudes of individual emitters. The general analysis is illustrated analytically for the case of the Gaussian approximation for emitting sources. The reduction of the interferometry radii and suppression of the Bose-Einstein correlation functions for small sources due to the uncertainty principle are found. There is the positive correlation between the source size and the intercept of the correlation function. The peculiarities of the non-femtoscopic correlations caused by minijets and fluctuations of the initial states of the systems formed in pp and e^+e^- collisions are analyzed also. The factorization property for the contributions of femtoscopic and non-femtoscopic correlations into complete correlation function is found.

PACS numbers: 13.85.Hd, 25.75.Gz

PACS: *24.10.Nz, 24.10.Pa, 25.75.-q, 25.75.Gz, 25.75.Ld.*

Keywords: *correlation femtoscopy, Hanbury Brown - Twiss intensity interferometry, proton-proton collisions, uncertainty principle, coherence, HBT correlations.*

Corresponding author: *Yu.M. Sinyukov, Bogolyubov Institute for Theoretical Physics, Kiev 03680, Metrolohichna 14b, Ukraine. E-mail: sinyukov@bitp.kiev.ua*

I. INTRODUCTION

The correlation femtoscopy, or intensity interferometry method, is the direct tool to measure the spatial and temporal scales of extremely small and short-lived systems created in particle and nuclear collisions with accuracy of 10^{-15} m and 10^{-23} sec correspondingly. The method [1–3], is grounded on the Bose-Einstein (BE) or Fermi-Dirac (FD) symmetric properties of the quantum states. It has a deep analogy with the intensity interferometry telescope that was proposed by Hanbury Brown and Twiss for measurements of angular sizes of remote stars [4]. In distinction on standard telescopic and microscopic techniques based on the registration of intensities of light or particles, e.g., electrons, coming from (or through) the object, this method deals with the correlation between intensities of the source radiation registered by two (many) spatially separated parts of devices such as telescopes, reflectors, particle detectors, etc. In fact, it measures the correlations between numbers of emitted identical particles detected in separated parts of the detector.

The femtoscopic space-time structure of the systems is typically represented in terms of the interferometry radii. They are result of the Gaussian fit of the correlation function defined as a ratio of the two- (identical) particle spectra to the product of the single-particle ones. In the pioneer papers [2, 3] the measured interferometry radii were interpreted as the geometrical sizes of the systems. Later on it was found [5–7] that for typical systems formed in experiments with heavy ions, the above geometrical interpretation needs to be generalized. The treatment of the interferometry radii as the homogeneity lengths [8, 9] in the systems and crucial suggestion as for femtoscopy scanning of the source radiation in different momentum bins bring the possibility to analyze different parts of the source and explain the behavior of the interferometry radii. In addition, the practical method how to use the final state interactions (FSI) and effects of long-lived resonances to extract the BE correlations in relatively large systems created in heavy ion collisions has been proposed [10].

An another challenge, which is still actual, concerns the femtoscopy analysis of relatively small systems created in particle interactions such as pp and e^+e^- where the observed femtoscopic scales are approximately 1 fm or smaller. Typically, the suppression of the correlation function is fairly large in these processes. Here we will analyze the femtoscopy of such small systems accounting for the uncertainty principle, coherence of the radiation from spatially very closely set emitters and non-femtoscopic (non BE, FD and FSI) correlations. The latter appear due to the energy-momentum conservation law and incoherent contributions to the two- and single-particle spectra induced by particles clusterization in momentum space and fluctuations of initial conditions of the collisions processes. The detailed analysis of these theoretical problems can help to provide the correct femtoscopy study of the small systems.

II. THE BASIC IDEAS OF INTENSITY INTERFEROMETRY TELESCOPE.

The intensity interferometry method for the stars angular sizes measurement was proposed and realized first by Hanbury Brown and Twiss in [4] at the end of fifties of XX century. The electromagnetic radiation from the star is the mixture of different (partially) monochromatic wave trains which are mutually incoherent at the moment of radiation. To see the principal aspects let us consider the emission from the different sites of a radiating object. If the stellar object is close to observer, like our Sun, then one can easily select, say, the opposite sites (edges) of it. If the two telescopes are directed to those different sites and one measures the correlations between photon numbers coming to each of the two telescopes, then the waves from different sites of the source do not mix in the telescopes and the correlations between them are absent – the signals coming to the two telescope reflectors are mutually incoherent. If, however, the stellar object – let us consider now the double star system – is very remote, so that it is impossible to select only one of the two star by a telescope, the light from both stars will come to each of the two reflectors and become mutually (partially) coherent. For simplicity let us imagine that at some moment of time the telescopes register only one wave train from each star, and both these trains are equally polarized, and follow each other continuously. *Ipso facto* we ignore the real problems of the intensity interferometry method – how to extract the signal from noise, but preserve the principal point of this method.

A. The basic formalism

One can decompose the electric field strength into positive and negative frequency parts $\mathbf{E} = \mathbf{E}^{(+)}(\mathbf{x},t) + \mathbf{E}^{(-)}(\mathbf{x},t)$, $\mathbf{E}^{(-)}(\mathbf{x},t) = \left(E^{(+)}(\mathbf{x},t)\right)^{\dagger}$. The ideal photon counter reacts just to the product $\mathbf{E}^{(-)}(\mathbf{x},t)\mathbf{E}^{(+)}(\mathbf{x},t) = |\mathbf{E}^{(+)}(\mathbf{x},t)|^2$ [11]. Far from the stellar object the light is described good by the plane waves, so that

$$\mathbf{E}^{(+)}(\mathbf{x},t) = Ae^{i(\mathbf{k}_1\cdot\mathbf{x}-\omega_1 t)} + Be^{i(\mathbf{k}_2\cdot\mathbf{x}-\omega_2 t)} \tag{1}$$

The complex amplitudes A and B have stochastic independent phases: $A = |A(t)|e^{i\varphi_A(t,\mathbf{x})}$, $B = |B(t)|e^{i\varphi_B(t,\mathbf{x})}$ which are roughly constants during coherence time τ_{coh} of the wave train, e.g. $\tau_{coh} \approx 10^{-8}$ sec, so that being averaged over period of time $T >> \tau_{coh}$ the amplitudes and their product become zero: $\langle A\rangle = 0$, $\langle B\rangle = 0$, $\langle AB\rangle = 0$. The intensity of light, that is proportional to the number of photons registered by the telescope/reflector and photomultipliers at point $\mathbf{x}$, is

$$\begin{aligned}\langle I_{\mathbf{k}_1,\mathbf{k}_2}(x)\rangle &= \left\langle \mathbf{E}^2\right\rangle = \left\langle \mathbf{E}^{(-)}(\mathbf{x},t)\mathbf{E}^{(+)}(\mathbf{x},t)\right\rangle = \left\langle |\mathbf{E}^{(+)}(\mathbf{x},t)|^2\right\rangle \\ &= \left\langle |A|^2 + |B|^2 + AB^*e^{i((\mathbf{k}_1-\mathbf{k}_2)\cdot\mathbf{x}-(\omega_1-\omega_2)t)} + A^*Be^{-i((\mathbf{k}_1-\mathbf{k}_2)\cdot\mathbf{x}-(\omega_1-\omega_2)t)}\right\rangle \\ &= |A|^2 + |B|^2\end{aligned} \tag{2}$$

where we supposed that $|A(t)|, |B(t)|$ are almost independent on t.

The statistically averaged intensity registered by one of the telescopes contains the information only about (averaged) squared modulus of the amplitudes. However, the correlation of intensities C, defined as the ratio of the averaged product of intensities registered at the two space-time points $(\mathbf{x_1}, t_1)$ and $(\mathbf{x_2}, t_2)$, to the product of averaged intensities registered at these points, depends already on the differences of momenta and energies of the light quanta. To simplify notation we put $|A| = |B|$ and get

$$\begin{aligned}C &= \frac{\langle I_{\mathbf{k}_1,\mathbf{k}_2}(\mathbf{x}_1)I_{\mathbf{k}_1,\mathbf{k}_2}(\mathbf{x}_2)\rangle}{\langle I_{\mathbf{k}_1,\mathbf{k}_2}(\mathbf{x}_1)\rangle\,\langle I_{\mathbf{k}_1,\mathbf{k}_2}(\mathbf{x}_2)\rangle} = \frac{1}{4|A|^4}\left\langle |E^{(+)}(\mathbf{x_1},t)|^2|E^{(+)}(\mathbf{x_2},t)|^2\right\rangle \\ &= 1 + \frac{1}{2}\cos[(\mathbf{k}_1-\mathbf{k}_2)(\mathbf{x}_1-\mathbf{x}_2) - (\omega_1-\omega_2)(t_1-t_2)] \\ &\approx 1 + \frac{1}{2}\cos[\theta\,|\mathbf{k}|\,d + (x_{L1} - x_{L2} - t_1 + t_2)(\omega_1-\omega_2)]\end{aligned} \tag{3}$$

where one took into account that $\langle |A|^2 A^* B = 0\rangle$, etc. Here the $\theta = D/L$ is the angular size of the double star system with "transverse" distance D between stars in the plane perpendicular to the direction to the system, L is the distance to the system, d is the "transverse" distance between two telescopes, $|\mathbf{k}|$ is mean detected wave number, x_{iL} is the "longitudinal", directed to the system, coordinate of the i-telescope and $t_1 - t_2 \equiv \Delta$ is the signal delay between points $\mathbf{x_1}$ and $\mathbf{x_2}$. It is worthy of noting that the time resolution τ_{res} in the method has to be smaller than the coherence time, $\tau_{res} < \tau_{coh}$, in order to provide correlation measurement of photon numbers during the mutual coherence time τ_{coh} of the electromagnetic waves in points $\mathbf{x_1}$ and $\mathbf{x_2}$. For stationary process the averaging over large period of time T plays role of the averaging over the ensemble of events with duration time τ_{coh}. Also note that the condition of validity of formula (3) is

$$\frac{x_{2L} - x_{1L}}{c} - (t_2 - t_1) < \tau_{coh}, \tag{4}$$

Otherwise $\langle A(\mathbf{x_1}, t_1) A(\mathbf{x_2}, t_2)\rangle = 0, \langle B(\mathbf{x_1}, t_1) B(\mathbf{x_2}, t_2)\rangle = 0$ and the correlations disappear, $C = 1$, because of mutual incoherence of waves coming to the two telescopes.

B. The nature of HBT effect

The formula (3) demonstrates the idea how to measure the differences in momenta (and energies) of photons radiated by remote stars by measuring the correlation function depending on distances between telescopes/reflectors and time delay. In fact, the momentum difference in transverse plane is connected with angular size of a stellar object. As for the difference in the energy of photons, it is possible, in principle, to measure it with restriction given by Eq. (4), and this difference would be associated with different temperatures of the two stars, if such a situation could take place. However there is no direct connection of the difference in energy of radiated photons with the time and space scales of the stellar system and, therefore, only angular size of this object can be extracted in this way. The latter is the basic application of the intensity interferometry telescope, and the method was used to measure the angular sizes of single remote stars [1].

---

[1] In this case one deals not with two emitters, but with many emitters N at the whole stellar disk, and formula (3) is modified at $t_1 = t_2$ as follows: $C - 1 \propto \frac{1}{N^2} \sum_{j \neq i = 1}^{N} \exp\left[i(\mathbf{k}_{Ti} - \mathbf{k}_{Tj})(\mathbf{x}_{T1} - \mathbf{x}_{T2})\right] \overset{N \to \infty}{\rightarrow} (\frac{2J_1(\theta|\mathbf{k}|d)}{\theta|\mathbf{k}|d})^2$. Determining the distance corresponding to the first zero of the correlation function in this representation Hanbury Brown and Twiss have measured the angular size of Sirius and some other stars.

Despite the classical description of Hanbury Brown and Twiss (HBT) method, the detailed analysis of the measurements accounting for the principle of photon registration pointed to the quantum nature of the effect [11]. The relationship between classical and quantum descriptions of the electromagnetic waves has been established through the formalism of coherent states [11]. Appealing to quantum nature of the electromagnetic fields, one can say that the method of the measurement of the star angular size is based on the positive correlations between numbers of photons registered in two close *space-time* points because of Bose-Einstein statistics for these quanta.

If a double star system is so far from detectors that the latter can register only a few photons per τ_{res}, then one has to use the amplitudes of registration at points x_1 and x_2 of the two photons emitted with mutually random phases from the two stars a and b: $A(x_1) = \langle a|x_1\rangle + \langle b|x_1\rangle$ and $A(x_2) = \langle a|x_2\rangle + \langle b|x_2\rangle$. Then taking the ratio of the averaged modulus squared of symmetric (over photons permutation) two-photon amplitude $A(x_1, x_2) = A(x_1)A(x_2)$ to the product of the averaged single-photon amplitudes one can get again the correlation structure (3). Note that describing a registration of the two neutrino from the star one should use antisymmetrized amplitude $A(x_1, x_2) = \langle a|x_1\rangle \langle b|x_2\rangle - \langle a|x_2\rangle \langle b|x_1\rangle$.

III. THE BASIC IDEAS OF THE CORRELATION INTERFEROMETRY

In particle physics the positive correlations between numbers of identical pions with close momenta emitted from an interaction region in proton-antiproton annihilations were found in 1960 by Goldhaber et al. (GGLP effect) [1]. It was understood that the nature of the effect lies in quantum statistics for identical particles demanding the symmetrization of bosonic wave function. Later on, basing on this fundamental principle, Kopylov and Podgoretsky [2] developed the method of pion interferometry microscope, or correlation femtoscopy. They found an analogy between the correlation interferometry and Hanbury Brown/Twiss (HBT) stellar intensity interferometry [4]. As one can see below, the mathematical structure of the principal aspects of the two methods can be presented in identical form. However, physically the methods are "orthogonal": HBT method measures differences of momenta and energies of photons radiated by stellar objects while the correlation femtoscopy measures the space-time separation of the emission points. Correspondingly, the former is based on the analysis of particles correlations as they depend on the space-time separation of the telescopes/reflectors, while the latter deals with dependence of the correlations on the particles momenta differences. In this sense the term "HBT radii" that one often uses to present the femtoscopy measurements is not quite correct, as R. Lednický stresses [12].

A. Standard approach

The basic ideas of the correlation femtoscopy are described in many publications, e.g., Ref. [13]. We reproduce them here with, however, some important remarks. Let us suppose that two identical bosons (e.g., pions) are emitted from the two space-time points, $x_1 = (t_1, \mathbf{x}_1)$ and $x_2 = (t_2, \mathbf{x}_2)$ and then *propagate freely.* The wave function of a single particle at the initial time t_i in the configuration representation is $\delta^3(\mathbf{x} - \mathbf{x_i})$. At some time t in the *momentum* $p = (p^0 = E, \mathbf{p})$-representation it is $\psi_{x_i}(p,t) = \frac{1}{(2\pi)^{3/2}} e^{-iEt} e^{-i\mathbf{x}_i\mathbf{p}}$.

In the momentum representation (p_1, p_2) the two-boson wave function is symmetrized and has the form

$$\psi_{x_1,x_2}(p_1,p_2;t) = \frac{1}{\sqrt{2}(2\pi)^3}\left[e^{i\mathbf{p}_1\mathbf{x}_1}e^{i\mathbf{p}_2\mathbf{x}_2} + e^{i\mathbf{p}_2\mathbf{x}_1}e^{i\mathbf{p}_1\mathbf{x}_2}\right]e^{-i(E_1+E_2)t}. \tag{5}$$

Then the probability to find the two pions with momenta p_1, p_2 is expressed through the scalar product of differences of 4-momenta and 4-coordinates of the two-pion emission:

$$W_{x_1,x_2}(p_1,p_2) = |\psi_{x_1,x_2}(p_1,p_2;t)|^2 \propto 1 + \cos\left[(p_1 - p_2)\cdot(x_1 - x_2)\right]. \tag{6}$$

Comparing Eq.(3) with result (6), related to the interferometry telescope, one can see that they differ from each other by the factor 1/2 before the cosine. The attentive reader can notice the difference between the two cases. In the case of emission from a double star system the correlation of intensities accounts for all the possibilities: this is the correlation between intensities of the wave trains coming from different stars as well as from the same star. In opposite, when formula (6) is derived, it is supposed that if one boson (with momentum $\mathbf{p}_1$) is emitted from the point $\mathbf{x}_1$, then the another boson (with momentum $\mathbf{p}_2$) is emitted from different point $\mathbf{x}_2$ and vice versa, but the possibility when the two bosons are emitted from the same point, $\mathbf{x}_1$ or $\mathbf{x}_2$, is excluded. In the case of *independent* particles radiation all the possibilities have to be taken into account. For such a case one usually demonstrates the idea of the correlation femtoscopy method by means of factorization of the two-particle normalized emission function, $\rho(x_1,x_2)=\rho(x_1)\rho(x_2)$ and integrates the two-point probability (6) over space-time region. Then for the correlation function $C(p_1,p_2)$ one has (we ignore here possible correlations between coordinates and momenta of the emitted particles)

$$C(p_1,p_2)=\frac{W(p_1,p_2)}{W(p_1)W(p_2)}=\frac{1}{W(p_1)W(p_2)}\int d^4x_1d^4x_2\rho(x_1)\rho(x_2)W_{x_1,x_2}(p_1,p_2)$$
$$=1+\frac{1}{W(p_1)W(p_2)}\left|\int d^4x\rho(x)e^{iq\cdot x}\right|^2 \qquad (7)$$

where $q=p_1-p_2$. So, the probability to find the particles with momenta p_1, p_2 at very large times $t\to\infty$ is expressed through the Fourier image of the emission function. This is the typical basis of the correlation interferometry method allowing one to analyze size and shape of small systems. If ρ is the Gaussian-like emission probability that in some reference frame has the form $\rho(x)\propto\exp\left[-\sum_{i=1}^{3}\frac{x_i^2}{2R_i^2}\right]\delta(t-t_0)$, then (7) reads as

$$C(p_1,p_2)=1+\exp\left[-\sum_{i=1}^{3}q_i^2R_i^2\right]. \qquad (8)$$

For example, the typical resolving width of the correlation function $\overline{q}=50$ MeV corresponds to the size $4\cdot10^{-15}$ m = 4 Fm (Fermi) = 4 fm (femtometer). The origination of the name of the method – correlation femtoscopy is obvious from such estimates.

B. Correlation femtoscopy formalism under microscope

Problem however appears when we apply the basic formula (7) to the system with small number of emitters. For example, if there are the only two different emitting points: $\rho(x)=\frac{1}{2}(\delta^4(x-x_1)+\delta^4(x-x_2))$, then from (7) follows ($\rho_i=1/2$)

$$W_{x_1,x_2}(p_1,p_2)\propto\sum_{i,j=1,2}\rho_i\rho_j(1+\cos[(p_i-p_j)(x_i-x_j)])=1+\cos^2\left[\frac{1}{2}(p_1-p_2)(x_1-x_2)\right] \qquad (9)$$

The result is incorrect as we shall show.

To analyze the situation in detail let us consider single particle radiation from the two points. If the emission from the point x_1 is associated with the quantum state that is distinguishable from the state corresponding to emission from the point x_2, then the two amplitudes A_i are allowed to be realized with some probabilities ρ_i and one can build the corresponding diagonal density matrix:

$$\rho_1\colon\quad A_1(p) = e^{i\mathbf{p}\mathbf{x}_1}e^{-iEt} \quad \text{and} \quad \rho_2\colon\quad A_2(p) = e^{i\mathbf{p}\mathbf{x}_2}e^{-iEt};\quad \rho_1+\rho_2=1. \tag{10}$$

Here and below we omit multiplier $(2\pi)^{-3/2}$ since these factors cancel in the correlation function (see below). Note that $W_{x_1,x_2}(p) = \rho_1 A_1(p)A_1^*(p) + \rho_2 A_2(p)A_2^*(p) = 1$. In the case of the two-boson *independent* radiation from the two points there are the three different final states (amplitudes) A_{ij} that are realized with the corresponding probabilities ρ_{ij}:

$$\begin{gathered}\rho_{11}\colon\; A_{11}(p_1,p_2) = e^{i\mathbf{p}_1\mathbf{x}_1}e^{i\mathbf{p}_2\mathbf{x}_1}e^{-i(E_1+E_2)t};\;\; \rho_{22}\colon\; A_{22}(p_1,p_2) = e^{i\mathbf{p}_1\mathbf{x}_2}e^{i\mathbf{p}_2\mathbf{x}_2}e^{-i(E_1+E_2)t};\\ \rho_{12}\colon\quad A_{12}(p_1,p_2) = \frac{1}{\sqrt{2}}\left(e^{i\mathbf{p}_1\mathbf{x}_1}e^{i\mathbf{p}_2\mathbf{x}_2} + e^{i\mathbf{p}_1\mathbf{x}_2}e^{i\mathbf{p}_2\mathbf{x}_1}\right)e^{-i(E_1+E_2)t}\;;\;\; \rho_{11}+\rho_{22}+\rho_{12}=1.\end{gathered} \tag{11}$$

If $\rho_{ii}=\rho_i^2$ and $\rho_1=\rho_2=1/2$ then $\rho_{11}=\rho_{22}=1/4$ and $\rho_{12}=1/2$. Then the probability to find the two particles with momenta p_1, p_2 that are emitted independently from the two points x_1 and x_2 is

$$C(p_1,p_2) = \frac{W_{x_1,x_2}(p_1,p_2)}{W_{x_1,x_2}(p_1)W_{x_1,x_2}(p_2)} = \sum_{i\leqslant j=1,2}\rho_{ij}\left|A_{ij}\right|^2 = 1+\frac{1}{2}\cos\left[(p_1-p_2)(x_1-x_2)\right]. \tag{12}$$

The last result coincides with Eq. (3) for stellar intensity interferometry telescope. The result (12) takes into account that the emission of both quanta from the same point (say, x_1) brings no interference while the wrong account for the interference leads to the formula (7). The latter becomes true only for large number of *independent* emitters. Therefore the simplest "derivation" of the intensity interferometry method (7) has to be corrected to exclude the double accounting in the correlation function for the contribution associated with the particles pairs emitted from very close points. This correction is significant if the number of *independent* emitters is not large. We shall consider this effect in detail in the next section.

If the emission from points x_1 and x_2 is not independent at all (full coherence), then one cannot use the diagonal matrix density associated with the probabilities ρ_1,ρ_2 but only the pure state: $A_{x_1,x_2}(p) = A_1(p)+A_2(p)$. The amplitude of the two identical bosons symmetrized over p_1,p_2 in this case is: $A_{x_1,x_2}(p_1,p_2) = A_{x_1,x_2}(p_1)A_{x_1,x_2}(p_2)$ and

$$C(p_1,p_2) = \frac{W_{x_1,x_2}(p_1,p_2)}{W_{x_1,x_2}(p_1)W_{x_1,x_2}(p_2)} = 1. \tag{13}$$

C. Correlation femtoscopy in random phase representation

Both these cases of completely independent and fully coherent radiations from points x_1 and x_2 can be reproduced in the formalism of partially coherent phases [14]. To demonstrate it let us consider the two-point source with coordinates x_1 and x_2 and some undetermined phases $\phi(x_i)$ $(i = 1, 2)$ and express amplitude of single boson emission with momentum p

$$A_{x_1,x_2}(p) = (e^{i\mathbf{p}\mathbf{x_1}}e^{i\phi(x_1)} + e^{i\mathbf{p}\mathbf{x_2}}e^{i\phi(x_2)})e^{-iEt}. \tag{14}$$

The symmetrized amplitude of the two-boson radiation is

$$A_{x_1,x_2}(p_1, p_2) = A_{x_1,x_2}(p_1)A_{x_1,x_2}(p_2). \tag{15}$$

The probability to register the two identical particles with momenta p_1 and p_2 is

$$\overline{W(p_1, p_2)} =< A_{x_1,x_2}(p_1)A_{x_1,x_2}(p_2)A^*_{x_1,x_2}(p_1)A^*_{x_1,x_2}(p_2) > \tag{16}$$

where brackets mean the averaging over events with partially coherent phases (over all events of the emission). Then in the case of random phases, $\left\langle e^{i(\phi(x_2)-\phi(x_1))}\right\rangle = \delta^4(x_1 - x_2)$, we get result (12) and for fully coherent radiation when $\left\langle e^{i(\phi(x_2)-\phi(x_1))}\right\rangle = 1$ we get (13).

In the case of independent emitters in points x_1, x_2 the above phase average can be presented in the form [14]:

$$G_{12} = \left\langle e^{i(\phi(x_2)-\phi(x_1))}\right\rangle = \delta^4(x_1 - x_2) = I_{12}\delta(t_1 - t_2) \tag{17}$$

where I_{12} is the overlap integral

$$I_{ij} = \left| \int d^3\mathbf{x}\psi_{x_i}(t, \mathbf{x})\psi^*_{x_j}(t, \mathbf{x}) \right| . \tag{18}$$

As it is demonstrated, an independent emission with fully random phases corresponds to the mixed state with probabilities ρ_i for i-states. Such a description is possible only if bosons are emitted *independently* from different points x_i in *distinguishable/orthogonal* quantum states. The latter requirement is satisfied in above case of the flat momentum spectrum for each emitter, $f(p) = W(p) = const$, since the initial quantum states $\delta^3(\mathbf{x} - \mathbf{x_i})$ taken in different points i are orthogonal.

D. Uncertainty principle and formalism of partially coherent phases

If the momentum spectrum $f(p)$ is essentially not flat, this corresponds to the wave packet characterized by its center x_i and some finite width. One can discriminate between the different states i and j only if they are approximately orthogonal: the overlap integral (18) is small, $I_{ij} \ll 1$. In other words, the distance between the centers of emitters has to be larger than the width of the emitted wave packets. Since the latter is the inverse of the variance Δp of the momentum spectra, so $(x_i - x_j)^2 \geq 1/\Delta p^2$. The latter expresses the uncertainty principle: one can discriminate the wave packet i without noticeable violation of the particle spectrum, if the measurement, that localizes the particle's position somewhere inside the sphere with the center x_i and the diameter not less than $1/\Delta p$, unambiguously points to the quantum state i, but not j. So, the distance $\Delta x = (x_i - x_j)$ should satisfy uncertainty principle in the form $\Delta x^2 \Delta p^2 > 1$. Then the system presented by the superposition of the quantum states A_i with phases $\phi(x_i)$ like in Eq. (14), after a phase averaging (17) is approximated well by the mixed state with diagonal density matrix ρ_{ij}.

In relativistic physics there is another uncertainty principle: the measurement of the particle's momentum p has accuracy depending on the duration of the measurement δt: $\delta p \sim 1/\delta t$ [16]. So one can measure the time of particle emission without noticeable violation of the momentum spectrum with accuracy not better than $1/\Delta p$ and, so, the δ-function in Eq. (17) has to be smeared when one deals with the wave packets. So, for normalized wave packets, one cannot use the random phase approximation if the distance in space and time between emitters is less than the width $1/\Delta p$ of the wave packet (in units $\hbar = c = 1$). For example, if $f(\mathbf{p}) = \frac{1}{(2\pi\Delta p^2)^{3/2}} e^{-\frac{\mathbf{p}^2}{2\Delta p^2}}$, then $I_{12} = e^{-\frac{\Delta p^2(\mathbf{x}_1-\mathbf{x}_2)^2}{2}}$ at $t_1 = t_2$ and $\delta(t_1 - t_2) \Longrightarrow G^t_{12} \to e^{-\frac{\Delta p^2(t_1-t_2)^2}{2}}$. The last term expresses uncertainty principle for momentum-time measurements.

Therefore, the fully chaotic phases, or diagonal density matrix, are possible only under some conditions which we discussed before. For normalized quantum states emitted from the points i and j the phase average can be written as

$$G_{ij} = \left\langle e^{i(\phi(x_i)-\phi(x_j))} \right\rangle = I_{ij} e^{-\frac{\Delta p^2(t_i-t_j)^2}{2}} \qquad (19)$$

where I_{ij} is the overlap integral (18), and Δp is a variance of the momentum spectrum of emitters.

In opposite case when $(x_i - x_j)^2 \ll 1/\Delta p^2$ the states are indistinguishable and overlap integral (19) $G_{ij} \approx 1$ at $t_1 \approx t_2$ since at $x_i = x_j$ it is just normalization. Then we come back to the fully coherent radiation (13) which takes place for very close emitters [15]. So, in both limited cases of chaotic and fully coherent emission Eq. (19) leads to physically obvious results, and so we will use Eq. (19) at any distances between the emitters as accounting for uncertainty principle. In this way we solve the problem of distinguishability (or indistinguishability) of the quantum states ψ_i associated with different emission points x_i: we always sum i-amplitudes, not i-probabilities, and average the resulting distributions/spectra over partially coherent phases in correspondence with the requirement of maximal possible distinguishability and independence of different i-states compatible with uncertainty principle for momentum & position and energy-momentum & time measurements.

E. Femtoscopic correlations in multiparticle systems

We discussed above the simplest situation with the only two emitted pions. Typically, the basic formalism of correlation interferometry in multipion systems for inclusive momentum spectra is similar [17]: one can use the same basic formula (7) with the substitutions: $W(p) \to p^0 \frac{d^3N}{d^3p}$, $W(p_1, p_2) \to p_1^0 p_2^0 \frac{d^6N}{d^3p_1 d^3p_2}$, $\rho(x) \to S(x, p \equiv \frac{p_1+p_2}{2})$. The correlation between the space-time position x of the emission point and momentum p of emitted particle, reflected in p-dependence of the emission function $S(x, p)$, leads to dependence of the interferometry radii in formula (8) on the mean momentum p of pion pair, $R_i \to R_i(p)$ [6]. The $x - p$ correlation appears due to fast expansion of the multiparticle systems created in high energy A+A (or even p+p) collisions. The radii $R_i(p)$ are associated then with homogeneity lengths in i-direction in inhomogeneous expanding systems [8]. So, one can carry the two-pion correlation results from the two-pion system to multiparticle systems, if applies the above substitutions. In fact, our quantum-mechanical consideration is related to the rest frame of the homogeneous emitting subsystem forming the spectra in a vicinity of some momentum p.

The another aspect of multiparticle emission is that the two-pion wave function may not be always factorized as a subsystem: $\Psi = |\pi^+\pi^+\rangle\, |X\rangle$, where state X is related to the residual part of the total system. The symmetrization/antisymmetrization procedure has to be applied to the total system wave function Ψ. However, it can be difficult to provide it if one supposes the distinguishability of the radiation points as in the standard correlation interferometry method of independent sources [1, 2]. For example, let us suppose that one pion with momentum p_1, $\pi^+(p_1)$, is the primary particle emitted from the fireball near the point x_1 and another pion $\pi^+(p_2)$ with the momentum p_2 is emitted together with particle X at the resonance decay, $Res \to \pi^+(p_2) + X(p_3)$ (say, $\rho^0 \to \pi^+(p_2) + \pi^-(p_3)$) near the point x_2. Such a system may not have $\pi(p_1) \leftrightarrow \pi(p_2)$ symmetry because

i) if at the $\pi(p_1) \leftrightarrow \pi(p_2)$ exchange the momentum p_3 of the particle X is preserved, then the momentum of emitted resonance is changed and gets the other value $p_1 + p_3$. This new resonance state has typically another probability to form as compared to the previous resonance state. (In extreme case it can even happen that the value $(p_1 + p_3)^2$ instead of $p^2_{Res} = (p_2 + p_3)^2$, excludes the resonance decay into the pair $\pi^+(p_1)$, $X(p_3)$ because of the kinematics). In general all that suppresses the interference effect [18];

ii) if the exchange of π^+ momenta is accompanied by the change of p_3 by the value $q = p_1 - p_2$, $p_3 \to p'_3 = p_3 + q$, then the two configurations $\left|\pi^+_{x_1}(p_1)\pi^+_{x_2}(p_2)X(p_3)\right\rangle$ and $\left|\pi^+_{x_1}(p_2)\pi^+_{x_2}(p_1)X(p'_3)\right\rangle$ become distinguishable and so cannot interfere;

iii) if at $\pi(p_1) \leftrightarrow \pi(p_2)$ the particles $X(p_3)$ together with $\pi^+(p_2)$ are emitted by the resonance Res from the fireball near point x_1 and primary pion $\pi^+(p_1)$ radiates from x_2, the requirement of distinguishability of the emission points may lead to principal distinguishability of the points of emission x_1 and x_2 of the particle X that excludes the pion interference effect.

So, the exact symmetry of the total system at $p_1 \leftrightarrow p_2$ exchange can be lost in similar situations and the two configurations $\left|\pi^+_{x_1}(p_1)\pi^+_{x_2}(p_2)\right\rangle$ and $\left|\pi^+_{x_1}(p_2)\pi^+_{x_2}(p_1)\right\rangle$ of the two-pion subsystem can interfere only partially. This has to be taken into account in the transport models used for high energy collisions.

The attempt to estimate the limits of applicability of the standard correlation femtoscopy is done in Ref. [19]. It is argued that when $\left|\pi^+_{x_1}(p_1)\pi^+_{x_2}(p_2)\right\rangle \leftrightarrow \left|\pi^+_{x_1}(p_2)\pi^+_{x_2}(p_1)\right\rangle$ then the classical phase-space position of the residual part of the system is changed by the value $q(x_1 - x_2)$ along each direction i. If this change exceeds the size $2\pi\hbar$ of the elementary cell of the phase-space per one degree of freedom, then the *residual* system moves to another quantum state, and it can be, in principle, measured. Therefore the pion interference should disappear when $q_i R_i > 2\pi$ ($\hbar = c = 1$), where R_i is the effective system size/homogeneity length in i-direction. However this argumentation fails already in the simplest case when the system contains only identical particles. Such a system can be symmetrized with distinguishable and independent radiation points, and there is no principal possibility to measure the exchange of the momenta between two identical pions at any value of q using for this aim the residual system containing the same sort of pions. We think that the real picture of the pion interference can be restored on the basis of symmetrized/antisymmetrized amplitude of the *total* system with partially coherent phases in the way described in subsection IID.

IV. THE CORRELATION FEMTOSCOPY IN GAUSSIAN APPROXIMATION

Here we apply the basic ideas discussed in the previous section to construct the simple analytical model accounting for uncertainty principle in the correlation femtoscopy. We use the non-relativistic approximation in the rest frame of the source which has Gaussian sizes corresponding to the homogeneity lengths R_i in the corresponding part of the total expanding system. The Lorentz transformation of the results to a global system where source moves with 4-velocity u^{μ} is trivial enough. Let us introduce a single bosonic state $\psi_x(p,t)$ corresponding to emission of a single particle with mass m and 4-momentum p from 4-point x and its free propagation:

$$\psi_x(p,t) = e^{i(\mathbf{px}-p^0 t)} e^{i\varphi(x)} \tilde{f}(\mathbf{p}) \tag{20}$$

where $\varphi(x)$ is some phase and $\tilde{f}$ defines the primary momentum spectrum $f(\mathbf{p})$ that we take in the Gaussian form with variance $\Delta p = k$:

$$f(\mathbf{p}) = \tilde{f}^2(\mathbf{p}) = \frac{1}{(2\pi k^2)^{3/2}} e^{-\frac{\mathbf{p}^2}{2k^2}}. \tag{21}$$

The amplitude of the single-particle radiation from some 4-volume at very large times t_∞ can be written as a superposition of the wave functions $\psi_x(p)$ with some coefficients/distribution $\rho(x)$ for emission 4-points x. In the momentum representation it looks like

$$A(p,t) = c \int d^4x \psi_x(p,t)\rho(x) \tag{22}$$

where c is the normalization constant. Let us select the two directions: parallel to beam axis, it is marked as index "L" and orthogonal to it – transverse axis "T". The distribution function of the emission centers is supposed to be the Gaussian one with normalized to unity quantity $\rho^2(\mathbf{x}, t)$ which is the probability distribution in the case of random phases $\phi(x)$.

$$\rho(x) = \frac{1}{2\pi R_T \sqrt{R_L T}} e^{-\frac{\mathbf{x}_T^2}{4R_T^2} - \frac{x_L^2}{4R_L^2} - \frac{t^2}{4T^2}} \tag{23}$$

The single-particle momentum density averaged over events with different phase distributions is

$$\overline{W(p)} = c^2 \int d^4x d^4x' e^{ip(x-x')} \rho(x)\rho(x') \langle e^{i(\varphi(x)-\varphi(x'))} \rangle f(\mathbf{p}). \tag{24}$$

To calculate the phase average one needs first to calculate the overlap integral (18). In non-relativistic approach the wave function of the particle emitted from the point $\mathbf{x}_i$ at moment t_i in coordinate representation is at some moment t

$$\psi_{x_i}(t, \mathbf{r}) = \frac{1}{(2\pi)^{3/2}} \int f(\mathbf{p}) e^{-i\mathbf{p}(\mathbf{r}-\mathbf{x}_i)} e^{-i\frac{\mathbf{p}^2}{2m}(t_i - t)} d^3p. \tag{25}$$

Then the modulus of the overlap integral (18) is

$$I_{x_i x_j} = \left| \int d^3\mathbf{r} \psi_{x_i}(t, \mathbf{r}) \psi^*_{x_j}(t, \mathbf{r}) \right| = \frac{e^{-\frac{k^2(\mathbf{x}_i - \mathbf{x}_j)^2}{2(1+k^4(t_i-t_j)^2/m^2)}}}{\sqrt[4]{1 + k^4(t_i - t_j)^2/m^2}}. \tag{26}$$

To provide calculations in analytical form we substitute the squared time difference $(t_i - \ t_j)^2$ by the constant proportional to its mean value over emission region

$$(t_i - t_j)^2 \to a\langle (t_i - t_j)^2 \rangle = a \int d^4x_i d^4x_j \rho(x_i)\rho(x_j)(t_i - t_j)^2 = 4aT^2 \equiv \alpha T^2 \tag{27}$$

As we will demonstrate later, the results obtained within such a prescription reproduce with good accuracy the exact numerical calculations for momentum spectra and correlation functions with the values of α depending on the basic parameters of emission. Then the overlap integral (19) that accounts for uncertainty principle takes the form

$$G_{xx'} = \langle e^{i(\varphi(x)-\varphi(x'))} \rangle = \frac{e^{-\frac{k^2(\mathbf{x}-\mathbf{x}')^2}{2(1+\alpha k^4 T^2/m^2)}}}{\sqrt[4]{1 + \alpha k^4 T^2/m^2}} e^{-k^2(t-t')^2/2}. \tag{28}$$

Now one can obtain the one-particle spectrum (24). It is presented below for the case when homogeneity lengths are equal, $R_T = R_L = R$:

$$\overline{W(p)} = N e^{-\frac{\mathbf{p}^2}{2k^2} - \frac{2\mathbf{p}^2 R^2}{1+4k_0^2 R^2} - \frac{\mathbf{p}^4 T^2}{2m^2(1+4k^2T^2)}}, \tag{29}$$

where

$$k_0^2 = k^2/(1 + \alpha k^4 T^2/m^2). \tag{30}$$

The two-particle spectrum averaged over events with partially coherent phases is

$$\overline{W(p_1,p_2)} = c^4 \int d^4x_1 d^4x_2 d^4x_1' d^4x_2' e^{i(p_1x_1+p_2x_2-p_1x_1'-p_2x_2')} f(\mathbf{p}_1) f(\mathbf{p}_2)\rho(x_1)\rho(x_2)\rho(x_1')\rho(x_2') \cdot \\ \cdot \langle e^{i(\varphi(x_1)+\varphi(x_2)-\varphi(x_1')-\varphi(x_2'))}\rangle. \qquad (31)$$

The 4-points phase correlator supposing emitters to be chaotic and independent in a maximum possible way permitted by uncertainty principle is decomposed into the sum of products of the two-point correlators (19) and contains the three terms

$$\langle e^{i(\varphi(x_1)+\varphi(x_2)-\varphi(x_1')-\varphi(x_2'))}\rangle = G_{x_1x_1'}G_{x_2x_2'} + G_{x_1x_2'}G_{x_2x_1'} - G_{x_1x_2'}G_{x_2x_1'}G_{x_1x_2} \qquad (32)$$

where the third term removes the double accounting which appears in the sum of the first two terms when $x_1 \approx x_2 \approx x_1' \approx x_2'$. See discussion in the Subsection IIB.

Let us split transverse direction "T" into the two ones: *out* that is directed along the total transverse momentum of the pair $\mathbf{p}_{1T} + \mathbf{p}_{2T}$ and *side* that is orthogonal to the *out*-direction as well as to the longitudinal "L" axis. Then the calculations of the one- and two-particle spectra lead to the following correlation function in the variables of the half of the bosonic pair momenta sum $\mathbf{p} = \frac{\mathbf{p}_1+\mathbf{p}_2}{2}$ and the momenta difference $\mathbf{q} = \mathbf{p_1} - \mathbf{p_2}$

$$C(\mathbf{p},\mathbf{q}) = \frac{\overline{W(p_1,p_2)}}{\overline{\overline{W(p_1)W(p_2)}}} = 1 + e^{-q_T^2R_T^2\frac{4k_0^2R_T^2}{1+4k_0^2R_T^2} - q_L^2R_L^2\frac{4k_0^2R_L^2}{1+4k_0^2R_L^2} - \frac{(\mathbf{q}\cdot\mathbf{p})^2T^2}{m^2}\frac{4k^2T^2}{1+4k^2T^2}} - C_d(\mathbf{p},\mathbf{q}), \qquad (33)$$

where $\overline{W(p_i)}$ is defined by (24), k_0 is (30) and subtracted function C_d eliminates the double accounting at the averaging of the 4-points phase correlator, it is associated with the third term in Eq. (32). This term reduces the intercept of the correlation function as one can see below.

The coefficient α in the expression for k_0 should be chosen from the requirement of a good agreement between the correlation function calculated using the approximation (28) and the one calculated numerically using the exact expression for the space part of the phase correlator (26). For the typical freeze-out temperature $T_{f.o.} = m_\pi = m$ the value of k is $k = \sqrt{T_{f.o.}m} = 0.14$ GeV. Having fixed the value of this parameter we present in Fig.1 the comparison of the *side*-projections of the correlation functions calculated numerically using (26) without subtraction of the double accounting and corresponding analytical expressions (the first two terms in (33)). The values of α at different system sizes are presented.



FIG. 1. The comparison of the correlation functions in *side* direction without subtraction of the double accounting with corresponding analytical approximations. The α parameter values, which give a good agreement with the exact results, are presented for different system sizes. For R=1.5 and 5 fm the curves are merged, while for R=0.5 fm there is a disagreement with exact result at large q. The momentum dispersion $k = m = 0.14$ GeV, p=0, T=R.

For fairly small systems, $R \approx 0.5 - 1.5$ fm they are about unity, $\alpha \approx 1$ and are decreasing, $\alpha \to 0$, when the system size R grows up.

As it follows from Eq. (33) the observed Gaussian interferometry radii of the system are reduced as compared to the standard results (we mark it by *st* index) for the interferometry radii for the Gaussian source. The reductions for *side*- (S), *out*- (O) and *long*- (L) interferometry radii in the LCMS system ($p_L = 0$) are

$$\begin{aligned}
\frac{R_S^2}{R_{S,st}^2} &= \frac{4k_0^2 R_T^2}{1+4k_0^2 R_T^2} \\
\frac{R_O^2}{R_{O,st}^2} &= \left(R_T^2 \frac{4k_0^2 R_T^2}{1+4k_0^2 R_T^2} + T^2 v_{out}^2 \frac{4k^2 T^2}{1+4k^2 T^2}\right) / \left(R_T^2 + T^2 v_{out}^2\right) \\
\frac{R_L^2}{R_{L,st}^2} &= \frac{4k_0^2 R_L^2}{1+4k_0^2 R_L^2}
\end{aligned} \tag{34}$$

where $v_{out} = p_{out}/m \ll 1$.

By means of Eq. (31) one can calculate the subtracting correlation function C_d corresponding to the third term in Eq. (32). It brings the result

$$C_d(\mathbf{p},\mathbf{q}) = F(k_0^2R^2, k^2T^2)e^{-\frac{2\mathbf{q}^2k_0^2R^4\left(1+8k_0^2R^2\right)}{\left(1+4k_0^2R^2\right)\left(1+8k_0^2R^2+8k_0^4R^4\right)}-\frac{2k^2T^4(\mathbf{p}\cdot\mathbf{q})^2\left(1+8k^2T^2\right)}{m^2(1+4p^2T^2)(1+8k^2T^2+8k^4T^4)}} \tag{35}$$

$$F(k_0^2R^2, k^2T^2) = \sqrt{\frac{k_0}{k}}\frac{(1+4k^2T^2)^{1/2}\,(1+4k_0^2R^2)^{3/2}}{(1+8k^2T^2+8k^4T^4)^{1/2}\,(1+8k_0^2R^2+8k_0^4R^4)^{3/2}} \tag{36}$$

One can note that the pre-exponential coefficient $F(k_0^2R^2, k^2T^2)$ tends to zero at large $k_0^2R^2$ and/or k^2T^2 and tends to unity when both of these quantities tend to zero. In the former case of large sizes and/or hard radiation the intercept of the correlation function tends to 2 and in the latter case, when the sizes of the source are very small and/or radiation is soft, the intercept tends to 1. For example, at T=0

$$k^2R^2 \gg 1, \quad C(\mathbf{p},\mathbf{q}) = 1 + e^{-\mathbf{q}^2R^2} \tag{37}$$

which is associated with the standard results, and

$$k^2R^2 \ll 1, \quad C(\mathbf{p},\mathbf{q})) \approx 1 \tag{38}$$

The last result corresponds to undistinguishable positions of the emitters due to uncertainty principle in case of very small sources and/or soft particle radiation.

We illustrate realistic cases in Fig. 2 for *side*-projection of the correlation function. Despite the correlation function at small k^2R^2 looses the Gaussian form presented by the second term in (33), the Gaussian fits still follow the tendencies of the analytic results (34) which demonstrate the reduction of the interferometry radii when the sizes of the sources become small. The elimination of the double accounting leads also to the reduction of the intercept of the correlation function when the system size shrinks. So, there is the positive correlation between observed interferometry radii and the intercept when the system size is changing. The analytic results reproduce the exact ones at $\alpha = 2$ ($R = 0.1$ fm), $\alpha = 1.6$ ($R = 0.5$ fm), $\alpha = 0.8$ ($R = 1.5$ fm). It is worthy of noting that at large enough system sizes, $R > 3 - 4$ fm, the effects practically disappear.

V. NON-FEMTOSCOPIC CORRELATIONS

Considering the correlation interferometry of small systems we cannot pass by the phenomenon of the so-called non-femtoscopic correlations. For a relatively small systems the



FIG. 2. The behavior of the two-particle Bose-Einstein correlation function (*side*-projection) where the uncertainty principle and correction for double accounting are utilized. The momentum dispersion $k = m = 0.14$ GeV, p=0, T=R.

additional two-particle correlations may appear in the kinematic region typical for quantum statistical (QS) and FSI correlations. The well known example of such additional correlations is the correlation induced by the total energy and momentum conservation laws (see, e.g., Ref. [20]). As opposed to the QS and FSI correlations, which are familiar from the correlation femtoscopy method, and, so, are called sometimes as femtoscopy correlations, these correlations are not directly related to the spatio-temporal scales of the emitter, and are called, therefore, *non-femtoscopic* correlations. Since the latter affect noticeably the correlation functions for small systems, the interferometry radii extracted from complete correlation function in e. g. $p+p$ collisions depend on the assumption about the so-called correlation baseline – the strength and momentum dependence of the non-femtoscopic correlations [21–23]. It has an influence on an interpretation of the momentum dependence of the interferometry radii in $p+p$ collisions, where the possibility of hydrodynamic behavior of matter is questionable. Therefore for successful and unambiguous applications of the correlation femtoscopy method to elementary particle collisions one needs to know mechanisms of the non-femtoscopic correlations to separate the femtoscopic and non-femtoscopic correlations. The very important question is whether it is possible to factorize both these contributions to the the total correlation function.

A. Non-femtoscopic correlations from minijets

To study this problem we follow the ideas of the Ref. [24] and consider simple analytical model of three-particle emission accounting for non-femtoscopic correlations caused by minijets fragmentation and energy-momentum conservation law. Even if one ignores the QS correlations, the amplitude of the three-particle emission, accounting for minijet fragmentation, cannot be expressed by the simple product of single-particle amplitudes, but is modified and can be written as follows [24]

$$A(\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3) = A(\mathbf{p}_1)A(\mathbf{p}_2)A(\mathbf{p}_3) \rightarrow A(\mathbf{p}_1)A(\mathbf{p}_2)A(\mathbf{p}_3)Q(\mathbf{p}_2, \mathbf{p}_3) \tag{39}$$

where factor $Q(\mathbf{p}_i, \mathbf{p}_j) = \exp\left(-\frac{(\mathbf{p}_i - \mathbf{p}_j)^2}{\gamma^2}\right)$ describes the correlation between the particles with momenta $\mathbf{p}_i$ and $\mathbf{p}_j$ induced by the fact they came from the same minijet (or cluster in momentum space). Taking into account different possible cluster structures that can be formed by different pairs of particles one can write the single- and two-particle emission probabilities in the following way

$$W(\mathbf{p}_1) = \sum_i \frac{1}{3} \int dp_1^* dp_2^* dp_3^* \delta(\mathbf{p}_1 - \mathbf{p}_i^*) |A(\mathbf{p}_1^*, \mathbf{p}_2^*, \mathbf{p}_3^*) Q(\mathbf{p}_2^*, \mathbf{p}_3^*)|^2 \tag{40}$$

$$W(\mathbf{p}_1, \mathbf{p}_2) = \sum_{i \neq j} \frac{1}{6} \int dp_1^* dp_2^* dp_3^* \delta(\mathbf{p}_1 - \mathbf{p}_i^*) \delta(\mathbf{p}_2 - \mathbf{p}_j^*) |A(\mathbf{p}_1^*, \mathbf{p}_2^*, \mathbf{p}_3^*) Q(\mathbf{p}_2^*, \mathbf{p}_3^*)|^2 \tag{41}$$

And the corresponding correlation function is

$$C(\mathbf{p}, \mathbf{q}) = \frac{2}{3} C^{mix}(\mathbf{p}, \mathbf{q}) + \frac{1}{3} C^{jet}(\mathbf{p}, \mathbf{q}) \tag{42}$$

where $C^{mix}(\mathbf{p}, \mathbf{q})$ denotes the contribution to the correlation function corresponding to the pairs formed by the "independent" particle and the particle from the cluster while $C^{jet}(\mathbf{p}, \mathbf{q})$ is another contribution related to the pairs where both particles belong to the cluster.

To describe the correlations induced by the momentum conservation law one should include certain delta-functions to the expressions for $W(\mathbf{p})$ and $W(\mathbf{p}_1, \mathbf{p}_2)$. Here we use the "soft" form of the momentum conservation law where the delta-function is substituted by its gaussian-like approximation, keeping in mind that the conservation law being strictly kept for the whole system of produced particles can be only approximately fulfilled for the subsystem of identical bosons under consideration.

$$\delta(\mathbf{p}_1^* + \mathbf{p}_2^* + \mathbf{p}_3^*) \rightarrow \Delta(\mathbf{p}_1^*, \mathbf{p}_2^*) = C e^{-(\mathbf{p}_1^* + \mathbf{p}_2^* + \mathbf{p}_3^*)^2/d^2} \tag{43}$$



FIG. 3. The correlation function $C(q)$ accounting for the non-femtoscopic correlations due to minijet/cluster formation and momentum conservation law, $C_{cl}(q)$, is compared with the pure QS correlation function $C_{BE}(q)$. One can see that the full CF is quite well factorized into the QS and non-femtoscopic parts. The model parameters are $\gamma = 0.5$ GeV, $p = 0.35$ GeV/c, $k = 0.1$ GeV/c, $R = 1$ fm

Fig. 3 illustrates the relation between the femtoscopic and non-femtoscopic correlations in our simple model. One can see that for small systems with the size ~ 1 fm the full correlation function can be factorized in the two parts corresponding to the femtoscopic and non-femtoscopic correlations.

B. Non-femtoscopic correlations from fluctuating initial state

The another type of correlations not induced by the quantum statistics effects are the correlations connected with existing of subensembles of events with different emission functions that leads to the corresponding fluctuations in single-particle and two-particle momentum spectra. In hydrodynamical models of nucleus-nucleus and proton-proton collisions these fluctuations can be caused by asymmetrically fluctuating initial densities used for the hydro stage of the model.

Let us consider the effect of such correlations on the resulting correlation function again on the example of simple analytical model. The probabilities of the single- and two-particle



FIG. 4. The non-femtoscopic correlation function (”baseline”) at the fluctuating initial conditions for hydrodynamic expansion of the system. Depending on the relation between value of the parameter a, characterizing the strength of the fluctuations and the parameter $d = 1$ GeV, that describes the softening of the momentum conservation law for the pion subsystem, the baselines can be growing, constant or decreasing with momentum difference q.



FIG. 5. Factorization of the femtoscopic and non-femtoscopic correlations when the latter are induced by fluctuating initial conditions for hydrodynamics

emission in case of fluctuating emission function should be averaged over the ensemble of states corresponding to the events with different initial conditions

$$W(p) = \sum_i \rho(u_i) W(p; u_i) \tag{44}$$

$$W(p_1, p_2) = \sum_i \rho(u_i) W_i(p_1, p_2; u_i), \tag{45}$$

where u_i denotes the i^{th}-type of initial conditions and $\rho(u_i)$ is the distribution over initial conditions, $\sum_i \rho(u_i) = 1$. The corresponding expression for the non-femtoscopic correlation function takes the form

$$C_{NF}(p_1, p_2) = \frac{\sum_i \rho(u_i) W(p_1; u_i) W(p_2; u_i)}{\sum_i \rho(u_i) W(p_1; u_i) \sum_j \rho(u_j) W(p_2; u_j)} \tag{46}$$

Let us consider the simple model where the single-particle u_i-dependent distribution $W(p_1; u_i)$ are obtained using the modified momentum spectra $f(\mathbf{p}) \propto e^{-\frac{\mathbf{p}^2}{2k^2}} \rightarrow e^{-\frac{(\mathbf{p}-\mathbf{u})^2}{2k^2}}$, where $\rho(u_i)$ distribution is supposed to be the Gaussian $\rho(\mathbf{u}) = \frac{a^3}{\pi^{3/2}} e^{-\mathbf{u}^2 a^2}$. The momentum conservation law we use, as in the previous subsection, in the soft form using the factor $\Delta(p_1, p_2) = C e^{-(p_1+p_2+\ldots)^2/d^2}$.

Depending on the ratio between parameters d describing momentum conservation and a describing initial conditions fluctuations the baseline formed by the C_{NF} function can be growing, constant or decreasing with q (see Fig. 4).

We see that fluctuating initial conditions for hydro can cause non-femtoscopic correlations similar to those resulting from minijet fragmentation. Fig. 5 illustrates the factorization of its contribution to the full correlation function.

VI. CONCLUSIONS

We discussed the principal problems of the correlation interferometry of small sources, in particular, the similarity and the principal difference between Hanbury Brown & Twiss intensity interferometry and pion correlation interferometry methods. The main subject of the paper is the effects caused by not complete distinguishability of the emitting points in the source because of the uncertainty principle. To account for the latter the formalism of partially coherent phases in the amplitudes of individual emitters is developed. The specific treatment is required for elimination of the double accounting to the correlation function from the configurations in four-points phase correlator when all individual emission centers are close. It is shown that the effects are significant for the systems with the sizes and emission duration times about 1 fm, and they are expressed in the reduction of the interferometry radii and suppression of the Bose-Einstein correlation functions. There is the positive correlation between the source size and the intercept of the correlation function. At the typical slopes of the momentum spectra in $p+p$ and $A+A$ collisions these effects are negligible for fairly large sources with radii about or more than 3-4 fm.

The peculiarities of the non-femtoscopic correlations caused by minijets and fluctuations of the initial states of the systems formed in pp and e^+e^- collisions are analyzed also. The factorization property for the contributions of femtoscopic and non-femtoscopic correlations into complete correlation function is demonstrated in the simple analytic models.

The application of the theoretical results to pp and e^+e^- collisions will be done in the separate work.

VII. ACKNOWLEDGMENT

The authors are grateful to R. Lednicky for fruitful discussions. The research was carried out within the scope of the EUREA: European Ultra Relativistic Energies Agreement (European Research Group: "Heavy ions at ultrarelativistic energies"), and is supported by the National Academy of Sciences of Ukraine (Agreement F3/2012) and by the State fund for fundamental researches of Ukraine (Agreement F33/42-2012).

---

[1] G.Goldhaber, S.Goldhaber, W. Lee, A. Pais//Phys. Rev. **120** (1960) 325.

[2] G.I. Kopylov, M.I. Podgoretsky, Sov. J. Nucl. Phys.: **15** (1972) 219; **15** (1972) 219 **18** (1973) 336; **19** (1974) 215.

[3] G. Cocconi, Phys. Lett. **B49** (1974) 459.

[4] R. Hanbury Brown, R.Q. Twiss, Nature, **177** (1956) 27;
R. Hanbury Brown, R.Q. Twiss, Proc. Roy. Soc. A **242** (1957) 300; **243** (1957) 291.

[5] S. Pratt, Phys. Rev. D **33** (1986)1314.

[6] A.N. Makhlin, Yu.M. Sinyukov, Sov. J. Nucl. Phys. **46** (1987) 345;
A. N. Makhlin, Yu. M. Sinyukov, Z. Phys. C **39** (1988) 69;

Yu. M. Sinyukov, Nucl. Phys. A **498** (1989) 151.

[7] Y. Hama, S.S. Padula, Phys. Rev. D **37** (1988) 3237.

[8] Yu.M. Sinyukov, Nucl.Phys. A **566** (1994) 589c;
Yu.M. Sinyukov, in: Hot Hadronic Matter: Theory and Experiment, eds. J. Letessier, H.H. Gutbrod and J. Rafelski (Plenum, New York) 1995, 309.

[9] S.V. Akkelin, Yu.M. Sinyukov, Phys. Lett. B **356** (1995) 525.

[10] Yu.M.Sinyukov, R.Lednický, J.Pluta, B.Erazmus, S.V.Akkelin Phys. Lett B432 (1998) 248.

[11] R. Glauber in book "Quantum optics and electronics" (Eds. C. DeWitt, A. Blandin, C. Cohen-Tannoudji), Gordon and Breach, Scientific Publishers, New York - London - Paris, 1965.

[12] R. Lednický, Yad. Phys. **67** (2004) 73.

[13] M. Gyulassy, S.K. Kauffmann, L.W. Wilson, Phys. Rev. C **20** (1979) 2267.

[14] Yu.M.Sinyukov, A.Yu.Tolstykh. Z.Phys.C - Particles and Fields **61** (1994) 593.

[15] R. Lednický, V.L. Lyuboshits, M.I. Podgoretsky. J.Nucl Phys. **38** (1983) 251.

[16] V.V. Berestetsky, E.M. Lifshits, L.P. Pitaevsky, "Relativistic Quantum Theory" Part 1,"Nauka", Moscow, 1968.

[17] U. A. Wiedemann and U. Heinz, Phys. Rep. 319 (1999) 145.

[18] M.I. Podgoretsky, R. Lednický, Preprints JINR R2-82-181, R2-82-327, Dubna, 1982.

[19] A.I. Golohvastov, Yad. Phys. **65** (2002) 196.

[20] Z. Chajęcki, M. Lisa, Phys. Rev. C **78**, 064903 (2008); Phys. Rev. C **79** (2009) 034908 .

[21] K. Aamodt, *et al.* (ALICE Collaboration), Phys. Rev. D **82** (2010) 052001 .

[22] K. Aamodt, *et al.* (ALICE Collaboration), arXiv:1101.3665 [hep-ex].

[23] M.M. Aggarwal, *et al.* (STAR Collaboration), Phys. Rev. C **83**, 064905 (2011).

[24] S.V. Akkelin, Yu.M. Sinyukov, Phys.Rev.D 85 (2012) 074023.